






	Skagit County NFP	WA State NFP	National NFP	NFP Objective
Maternal smoking during pregnancy, relative change (N)	-14% (119)	-17% (3,894)	-16% (110,321)	-20%
Maternal alcohol use during pregnancy, relative change (N)	-100% (119)	-14% *** (3,957)	-20% *** (112,060)	-20%
Low-birth weight infants, Percent (N)	7% (162)	8% (5,311)	10% (137,249)	NA
Very low birth weight, Percent (N)	1% (162)	1% (5,311)	3% (137,249)	NA
Note: For Washington St	alysis of data from Nurse e cumulative from progr ate and National NFP c Igit County NFP) are inc	am initiation throug omparison groups,	h December 31, 20 statistically significa	14. Int differences

Capacity Levels	Measurement Domains
Coalition Capacity	Leadership and infrastructure
	Communications
Network Capacity	Goal-directed networks
	Community cross-sector partnerships
	Shared goals
Community-based Solutions	Community problem-solving processes
	Focus on equity
	Data use for improvement and accountability
Community-wide Impact	Multilevel strategies
	Diverse engagement and empowerment
	Scale of work



	Number of			
APPI Sites	Sample Total	responses	Response rate	
Overall	276	233	84.4%	
NCW	39	29	74.7%	
Okanogan	42	35	83.3%	
Skagit	52	42	80.8%	
Walla Walla	76	69	90.8%	
Whatcom	67	58	86.6%	

ARC ³ Domain	Scores and	Reliabilities
-------------------------	------------	---------------

2.80 (0.68) 2.79 (0.68) 2.44 (0.66)	0.63 - 0.84 0.65 - 0.68 0.58 - 0.66	0.82 0.78
. ,		
2.44 (0.66)	0.58 - 0.66	1
		0.76
2.43 (0.86)	0.70 – 0.86	0.87
2.70 (0.78)	0.64 – 0.78	0.81
2.95 (0.70)	0.71 – 0.77	0.76
2.47 (0.78)	0.66 – 0.80	0.79
2.97 (0.70)	0.64 - 0.86	0.84
2.41 (0.64)	0.72 – 0.87	0.85
2.22 (0.81)	0.66	0.79
	2.95 (0.70) 2.47 (0.78) 2.97 (0.70) 2.41 (0.64)	2.95 (0.70) 0.71 - 0.77 2.47 (0.78) 0.66 - 0.80 2.97 (0.70) 0.64 - 0.86 2.41 (0.64) 0.72 - 0.87

